

NAME OF THE DOCUMENTPATENT SPECIFICATIONTITLE OF THE INVENTION NEW GENES

RECEIVED

AUG 21 2002

DETAILED EXPLANATION OF THE INVENTION

OFFICE OF PATENTS

CLAIMS:

1. A protein selected from the group consisting of:
 - (a) a protein which consists of an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89; and
 - (b) a protein which is capable of activating NF- κ B and consists of an amino acid sequence having at least one amino acid deletion, substitution or addition in an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89.
2. A protein which is capable of activating NF- κ B and comprises an amino acid sequence having a homology of at least 50% to the amino acid sequence of the protein according to claim 1.
3. A gene which comprises a nucleotide sequence encoding a protein selected from the group consisting of:
 - (a) a protein which consists of an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89; and
 - (b) a protein which is capable of activating NF- κ B and consists

of an amino acid sequence having at least one amino acid deletion, substitution or addition in an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89.

4. A gene which comprises a DNA sequence selected from the group consisting of:

(a) a DNA sequence which consists of a nucleotide sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90; and

(b) a DNA sequence which encodes a protein capable of activating NF- κ B and hybridizes with the DNA sequence of above item (a) under stringent conditions.

5. A gene comprising a nucleotide sequence which encodes a protein capable of activating NF- κ B and has a homology of at least 50% to the nucleotide sequence of the gene according to claim 3 or 4.

6. A protein encoded by a gene according to any one of claims 3 to 5.

7. A recombinant vector which comprises a gene according to any one of claims 3 to 5.

8. A transformant which comprises a recombinant vector according to claim 7.

9. A process for producing a polypeptide, which comprises expressing the polypeptide encoded by the DNA from the transformant according to claim 8.

10. A method for diagnosing a disease associated with the expression or activity of NF- κ B in a subject, which comprises the steps of:

- (a) determining the presence or absence of a mutation in the NF- κ B coding nucleotide sequence in the genome of the subject; and/or
- (b) measuring the amount of the expression of the protein according to claim 6 in the subject.

11. A method for screening a compound which inhibits or activates NF- κ B activation, which comprises the steps of:

- (a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal;
- (b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell;
- (c) contacting the transformed cell with one or more candidate compounds; and
- (d) measuring the detectable signal.

12. A process for producing a substance which inhibits or activates NF- κ B activation, which comprises the steps of:

- (a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal;
- (b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell;
- (c) contacting the transformed cell with one or more candidate compounds;
- (d) measuring the detectable signal; and
- (e) identifying and isolating the candidate compound in which the signal of the inhibition or activation of NF- κ B activation is

detected.

13. A compound which inhibits or activates NF- κ B activation, which is obtained by the process according to claim 12.

14. A kit for screening a compound which inhibits or activates NF- κ B activation, which comprises:

(a) a host cell transformed with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal; and

(b) a medium which permits the expression of the gene in the host cell.

15. A monoclonal or polyclonal antibody directed against a protein according to claim 1 or 2.

16. A process for producing a monoclonal or polyclonal antibody according to claim 15, which comprises using a protein according to claim 1 or 2 as an antigen.

17. An antisense oligonucleotide against a gene according to any one of claims 3 to 5.

18. A ribozyme which inhibits NF- κ B activation.

19. A method for treating a disease, which comprises using a compound according to claim 13, and/or a monoclonal or polyclonal antibody according to claim 15, and/or an antisense oligonucleotide according to claim 17, and/or a ribozyme according to claim 18.

20. A pharmaceutical composition which comprises a compound according to claim 13 as an active ingredient.

21. A pharmaceutical composition which comprises a monoclonal or polyclonal antibody according to claim 15 as an active ingredient.

22. A pharmaceutical composition which comprises an antisense oligonucleotide according to claim 17 as an active ingredient.

23. The pharmaceutical composition according to claim 20, wherein

the target disease is selected from the group consisting of inflammation, autoimmune diseases, infections diseases and cancers.

24. A method for obtaining a new gene having a function, which comprises a method of using the oligo-capping method to construct a full-length cDNA library, and a method of using a reporter gene indicative of the presence of a protein having the function.

25. A computer-readable medium on which a sequence data set has been stored, said sequence data set comprising at least one nucleotide sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90, and/or at least one amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89.

26. A method for calculating a homology, which comprises comparing data on a medium according to claim 25 with data of other nucleotide sequences and/or amino acid sequences.

DETAILED EXPLANATION OF THE INVENTION

TECHNICAL FIELD OF THE INVENTION

The present invention relates to a protein capable of activating NF- κ B, a DNA sequence encoding the protein, a method for obtaining the DNA, a recombinant vector containing the DNA, a transformant containing the recombinant vector, and an antibody which specifically reacts with the protein. The present invention

also relates to use of the protein, DNA molecule or antibody of the invention in the diagnosis, treatment or prevention of diseases associated with the excessive activation or inhibition of NF- κ B.

The present invention also relates to a method for screening a substance capable of inhibiting or activating NF- κ B activation by using the protein, DNA, recombinant vector and transformant.

DESCRIPTION

The transcription factor NF- κ B (nuclear factor kappa B) plays an important role in the transcriptional regulation of various genes involved in inflammation and immunological reactions. NF- κ B is a homo- or heterodimer which belongs to the Rel family. In unstimulated conditions, NF- κ B normally resides in the cytoplasm as an inactive form by forming a complex with an I κ B inhibitory protein (Inhibitory protein of NF- κ B) to mask the nuclear transport signal of NF- κ B.

When cytokines such as interleukin (IL)-1 and tumor necrosis factor (TNF)- α stimulate cells, I κ B is phosphorylated by IKK (I κ B kinase) and degraded by the 26S proteasome through ubiquitination. The released NF- κ B moves to the nucleus, where it binds to the DNA sequence called the NF- κ B binding sequence and induces the transcription of the gene, which is under control of NF- κ B is believed to regulate the expression of genes such as those for immunoglobulins, inflammatory cytokines (e.g., IL-1 and TNF- α), interferons and cell adhesion factors. NF- κ B is involved in inflammation and immune responses through the expression induction of these genes.

The inhibition of the function or activation of NF- κ B may

inhibit the expression of many factors (proteins) involved in inflammatory or immunological diseases or other diseases such as tumor proliferation. Thus, NF- κ B is a promising target for medicaments against diseases caused or characterized by autoimmunity or inflammation [see e.g., Clinical Chemistry 45, 7-17 (1999); J Clin. Pharmacol. 38, 981-993 (1998); Gut 43, 856-860 (1998); The New England Journal of Medicine 366, 1066-1071 (1997); TIPS 46-50 (1997); The FASEB Journal 9, 899-909 (1995); Nature 395, 225-226 (1998); Science 278, 818-819 (1997); Cell 91, 299-302 (1997)].

Extracellular information is converted into a certain signal, which passes through the cell membrane and goes through the cytoplasm to the nucleus, where it regulates the expression of the target gene and causes cell responses. Therefore the elucidation of the mechanism of intracellular signal transduction from extracellular stimuli to NF- κ B activation is of very important significance, because it provides very important means of developing new medicaments or therapies against autoimmune diseases and diseases exhibiting inflammatory symptoms.

It is believed that the signal transduction pathway from certain cell stimulation to NF- κ B activation includes many steps mediated by various transmitters such as protein kinases. Therefore it is desirable for more efficient drug discovery to identify the transmitters which play a key role in the pathway, and to focus research on the transmitters to establish a new drug-screening method. Some signaling molecules involved in NF- κ B activation have been identified [e.g., IKK, ubiquitination enzymes and the 26S proteasome described above, as well as TNF receptor associated factor 2 (TRAF2) and NF- κ B inducing kinase (NIK)].

However, most of the mechanism of NF- κ B activation remains unknown, and it has been desired new signaling molecules to be identified and further the NF- κ B activation mechanism to be elucidated.

PROBLEM TO BE SOLVED BY THE INVENTION

The object of the present invention is to identify a new gene and protein capable of activating NF- κ B, and to provide a method of use of them in medicaments, diagnostics and therapy. That is, the present invention provides a new protein capable of activating NF- κ B, a DNA sequence encoding the protein, a recombinant vector containing the DNA, a transformant containing the recombinant vector, a process for producing the protein, an antibody directed against the protein or a peptide fragment thereof, and a process for producing the antibody.

The present invention also provides a method for screening a substance capable of inhibiting or activating NF- κ B activation, a kit for the screening, a substance capable of inhibiting or activating NF- κ B activation obtainable by the screening method or the screening kit, a process for producing the substance, a pharmaceutical composition containing a substance capable of inhibiting or activating NF- κ B activation, etc.

MEANS TO SOLVE THE PROBLEM

Recently, random analysis of cDNA molecules has been intensively carried out to analyze various genes which are expressed in vivo. The cDNA fragments thus obtained have been entered for databases and published as ESTs (Expressed Sequence Tags, e.g., <http://www.ncbi.nlm.nih.gov/dbEST>). However, ESTs are merely sequence information, and it is difficult to predict their functions. ESTs are also arranged in UniGene (<http://www.ncbi.nlm.nih.gov/UniGene>), and about 80,000 human genes

have been registered until now. However, most of these ESTs have their 5' end nucleotide sequences deleted, and contain no translation initiation site. Therefore it is unlikely that such analysis will directly lead to gene functional analysis such as the analysis of protein functions on the assumption of the determination of mRNA coding regions and the understanding of gene expression control by the analysis of promoters.

On the other hand, one method to elucidate functions of gene products (i.e., proteins) is transient expression cloning method using animal cells [see e.g., "Idenshi Kougaku Handbook (Genetic Engineering Handbook)", an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.]. This method involves transfecting animal cells with a cDNA library constructed using an animal cell expression vector to directly express a functional protein, and identifying and cloning the cDNA based on the biological activity of the protein having an effect on the cells. This method requires no chemical information (amino acid sequences and molecular weights) regarding the target protein product as a prerequisite, and allows the identification of cDNA clones by detecting specific biological activity of the protein expressed in the cells or culture.

For the efficient expression cloning, there is a need to devise a method of preparing a cDNA library. Several methods have been widely used to construct cDNA libraries [e.g., the method of Gubbler-Hoffman: Gene 25 (1983); and the method of Okayama-Berg: Mol. Cell. Biol. 2 (1982)]. However, most of the cDNA molecules prepared by these methods have their 5' end nucleotide sequences deleted, and thus these methods rarely produce full-length cDNA, a complete DNA copy of mRNA. This is because the reverse

transcriptase used to prepare cDNA from mRNA does not necessarily have high efficiency in producing full-length cDNA. Therefore it is necessary to improve these prior art methods in order to efficiently carry out the above expression cloning.

In addition, in order to carry out the functional analysis of genes, it is essential to clone full-length cDNA sequences and express proteins from them. Therefore, it has been necessary to construct cDNA libraries containing enriched full-length cDNA for efficient expression cloning.

The present inventors have intensively studied to solve the above problems. As a result, the present inventors have succeeded in constructing a full-length cDNA library by using the oligo-capping method; establishing an assay system using 293EBNA cells; and isolating a new DNA (cDNA) encoding a protein having a function of activating NF- κ B by using the assay system. This new DNA molecule induced NF- κ B activation by its expression in 293EBNA cells. This result shows that this new DNA is a signal transduction molecule involved in NF- κ B activation. Thus, the present invention has been completed.

That is, the present invention relates to:

- (1) A protein selected from the group consisting of:
 - (a) a protein which consists of an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89; and
 - (b) a protein which is capable of activating NF- κ B and consists of an amino acid sequence having at least one amino acid deletion, substitution or addition in an amino acid sequence selected from

the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89;

(2) A protein which is capable of activating NF- κ B and comprises an amino acid sequence having a homology of at least 50% to the amino acid sequence of the protein according to above item (1);

(3) A gene which consists of or comprises a nucleotide sequence encoding a protein selected from the group consisting of:

(a) a protein which consists of an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89; and

(b) a protein which is capable of activating NF- κ B and consists of an amino acid sequence having at least one amino acid deletion, substitution or addition in an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89,;

(4) A gene which consists of or comprises a DNA sequence selected from the group consisting of:

(a) a DNA sequence which consists of a nucleotide sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90; and

(b) a DNA sequence which encodes a protein capable of activating

NF- κ B and hybridizes with the DNA sequence of above item (a) under stringent conditions;

(5) A gene comprising a nucleotide sequence which encodes a protein capable of activating NF- κ B and has a homology of at least 50% to the nucleotide sequence of the gene according to above item (3) or (4);

(6) A protein encoded by a gene according to any one of above items (3) to (5);

(7) A recombinant vector which comprises a gene according to any one of above items (3) to (5);

(8) A transformant which comprises a recombinant vector according to above item (7);

(9) A process for producing a polypeptide, which comprises expressing the polypeptide encoded by the DNA from the transformant according to above item (8).

The present invention also relates to a diagnostic method for a disease associated with the activation or inhibition of NF- κ B, namely:

(10) A method for diagnosing a disease associated with the expression or activity of NF- κ B in a subject, which comprises the steps of:

- (a) determining the presence or absence of a mutation in the NF- κ B coding nucleotide sequence in the genome of the subject; and/or
- (b) measuring the amount of the expression of the protein according to above item (6) in the subject.

The present invention further relates to screening for and production of factors which regulate NF- κ B activity; namely:

(11) A method for screening a compound which inhibits or activates NF- κ B activation, which comprises the steps of:

(a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal;

(b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell;

(c) contacting the transformed cell with one or more candidate compounds; and

(d) measuring the detectable signal;

(12) A process for producing a substance which inhibits or activates NF- κ B activation, which comprises the steps of:

(a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal;

(b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell;

(c) contacting the transformed cell with one or more candidate compounds;

(d) measuring the detectable signal; and

(e) identifying and isolating the candidate compound in which the signal of the inhibition or activation of NF- κ B activation is detected;

(13) A compound which inhibits or activates NF- κ B activation, which is obtained by the process according to above item (12);

(14) A kit for screening for a compound which inhibits or activates NF- κ B activation, which comprises:

(a) a host cell transformed with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal; and

(b) a medium which permits the expression of the gene in the host

cell.

The present invention also relates to:

- (15) A monoclonal or polyclonal antibody directed against a protein according to above item (1) or (2);
- (16) A process for producing a monoclonal or polyclonal antibody according to above item (15), which comprises using a protein according to above item (1) or (2) as an antigen;
- (17) An antisense oligonucleotide against a gene according to any one of above items (3) to (5);
- (18) A ribozyme which inhibits NF- κ B activation.

The present invention further relates to therapy and a pharmaceutical composition involving these substances; namely:

- (19) A method for treating a disease, which comprises using a compound according to above item (13), and/or a monoclonal or polyclonal antibody according to above item (15), and/or an antisense oligonucleotide according to above item (17), and/or a ribozyme according to above item (18);
- (20) A pharmaceutical composition which comprises a compound according to above item (13) as an active ingredient;
- (21) A pharmaceutical composition which comprises a monoclonal or polyclonal antibody according to above item (15) as an active ingredient;
- (22) A pharmaceutical composition which comprises an antisense oligonucleotide according to above item (17) as an active ingredient;
- (23) The pharmaceutical composition according to above item (20), wherein the target disease is selected from the group consisting of inflammation, autoimmune diseases, infections diseases and cancers.

The present invention also relates to:

(24) A method for obtaining a new gene having a function, which comprises a method of using the oligo-capping method to construct a full-length cDNA library, and a method of using a reporter gene indicative of the presence of a protein having the function;

(25) A computer-readable medium on which a sequence data set has been stored, said sequence data set comprising at least one nucleotide sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90, and/or at least one amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89;

(26) A method for calculating a homology, which comprises comparing data on a medium according to above item (25) with data of other nucleotide sequences and/or amino acid sequences.

At first, in order to further clarify the basic feature of the present invention, the present invention is explained by following how the present invention is completed. In order to obtain a new gene having a function of activating NF- κ B, the following experiments were carried out as shown in the examples. First, using the oligo-capping method, a full-length cDNA was produced from mRNA prepared from normal human lung fibroblasts (purchased from Sanko Junyaku Co., Ltd.), and a full-length cDNA library was constructed in which the cDNA was inserted into the vector pME18S-FL3 (GenBank Accession AB009864). Next, the cDNA library was introduced into E. coli cells, and plasmid preparation was carried out per clone.

Then, the pNK κ B-Luc reporter plasmid (STRATAGENE) containing a DNA encoding luciferase and the above full-length cDNA plasmid were cotransfected into 293-EBNA cells (Invitrogen). After 24 or 48 hours of culture, luciferase activity was measured, and the plasmid with significantly increased luciferase activity compared to that of a control experiment was selected (the selected plasmid showed a 10-fold or more increase in luciferase activity compared to that of the control experiment), and the entire nucleotide sequence of the cDNA cloned into the plasmid was determined. The protein encoded by the cDNA thus obtained shows that this protein is a signal transduction molecule involved in NF- κ B activation.

The present invention is explained definitely below.

The present invention includes a protein selected from the group consisting of (a) a protein which consists of an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89; and (b) a protein which is capable of activating NF- κ B and consists of an amino acid sequence having at least one amino acid deletion, substitution or addition in an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89. The Examples section described below demonstrate that the protein consisting of an amino acid sequence of the above SEQ ID NOS: 1, etc., is capable of activating NF- κ B.

Those skilled in the art can prepare a protein having the same NF- κ B activating activity as the protein having an amino acid

sequence of SEQ ID NOS 1, etc by means of appropriate substitution of an amino acid substitution in the protein using the known methods. One such method involves using conventional mutagenesis procedures for the DNA encoding the protein. Another method is, for example, site-directed mutagenesis (e.g., Mutan-Super Express Km Kit from Takara Shuzo Co., Ltd.). Mutations of amino acids in proteins may also occur in nature. Thus, the present invention also includes a mutated protein which is capable of activating NF- κ B and which has at least one amino acid deletion, substitution or addition compared to the corresponding SEQ ID NO amino acid sequence. The number of mutations is preferably up to 10, more preferably up to 5, most preferably up to 3.

Examples of such amino acid substitutions include substitutions within the following groups: (glycine, alanine), (valine, isoleucine, leucine), (aspartic acid, glutamic acid), (asparagine, glutamine), (serine, threonine), (lysine, arginine) and (phenylalanine, tyrosine).

Based on the DNA sequences (e.g., SEQ ID NO: 2) encoding a protein consisting of an amino acid sequence of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 or 89, or fragments thereof, those skilled in the art can routinely isolate a DNA with a high homology to these DNA sequences by using hybridization techniques and the like, and obtain proteins having the same NF- κ B activating activity as the protein having of an amino acid sequence of SEQ ID NOS 1, etc. Thus, the present invention also includes a protein which is capable of activating NF- κ B and comprises an amino acid sequence having a high homology to the amino acid sequence of above SEQ ID

NOS 1, etc. "High homology" refers to an amino acid sequence having a homology of at least 50%, preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, at the amino acid level.

The protein of the present invention may be natural proteins derived from any human or animal cells or tissues, chemically synthesized proteins, or proteins obtained by genetic recombination techniques. The protein may or may not be subjected to post-translational modifications such as sugar chain addition or phosphorylation.

The present invention also includes a gene encoding the above protein of the present invention. Examples of nucleotide sequences encoding a proteins consisting of an amino acid sequence of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 or 89 include nucleotide sequences of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 or 90. The gene of the present invention includes DNA substantially comprising these entire nucleotide sequences or fragments thereof, as well as its corresponding RNA. The DNA includes cDNA, genomic DNA, and chemically synthesized DNA. In accordance with the degeneracy of the genetic code, at least one nucleotide in the nucleotide sequence encoding a protein consisting of an amino acid sequence of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 or 89 can be substituted with other nucleotides without altering the amino

acid sequence of the protein produced from the gene. Therefore the DNA sequences of the present invention also include nucleotide sequences altered by substitution based on the degeneracy of the genetic code. Such DNA sequences can be synthesized using known methods.

The DNA of the present invention includes a DNA which encodes a protein capable of activating NF- κ B and hybridizes under stringent conditions with the DNA sequence of the above nucleotide sequence of SEQ ID NOS: 2, etc. The stringent conditions are apparent to those skilled in the art, and can be easily attained in accordance with various laboratory manuals such as T. Maniatis et al., Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory 1982, 1989.

That is, "stringent conditions" refer to overnight incubation at 37°C in a hybridization solution containing 30% formamide, followed by washing (three times) in 2 x SSC, 0.1% SDS for 10 minutes at room temperature, then followed by washing (two times) in 0.2 x SSC, 0.1% SDS for 10 minutes at 37°C. Preferred stringent conditions are overnight incubation at 42°C in a hybridization solution containing 40% formamide, followed by washing (three times) in 2 x SSC, 0.1% SDS for 10 minutes at room temperature, then followed by washing (two times) in 0.2 x SSC, 0.1% SDS for 10 minutes at 42°C. More preferred stringent conditions are overnight incubation at 42°C in a hybridization solution containing 50% formamide, followed by washing (three times) in 2 x SSC, 0.1% SDS for 10 minutes at room temperature, followed by washing (two times) in 0.2 x SSC, 0.1% SDS for 10 minutes at 50°C. The DNA sequence thus obtained must encode a protein capable of activating NF- κ B.

The present invention also includes a gene comprising a nucleotide sequence which encodes a protein capable of activating NF- κ B and has a homology of at least 50% to the nucleotide sequence of the gene according to above item (3) or (4).

The homology of the nucleotide sequences is more preferably at least 70%, even more preferably at least 80%, most preferably at least 90%.

The above DNA sequence of the present invention can be used to produce the above protein using recombinant DNA techniques. In general, the DNA and peptide of the present invention can be obtained by:

- (A) cloning the DNA encoding the protein of the present invention;
- (B) inserting the DNA encoding the entire coding region of the protein or a part thereof into an expression vector to construct a recombinant vector;
- (C) transforming host cells with the recombinant vector thus constructed; and
- (D) culturing the obtained cells to express the protein or its analogue, and then purifying it by column chromatography.

General procedures necessary to handle DNA and recombinant host cells (e.g., *E. coli*) in the above steps are well known to those skilled in the art, and can be easily carried out in accordance with various laboratory manuals such as T. Maniatis et al., *supra*. All the enzymes, reagents, etc., used in these procedures are commercially available, and unless otherwise stated, such commercially available products can be used according to the use conditions specified by the manufactures' instructions to attain completely its objects. The above steps (A) to (D) can be further illustrated in more details as follows.

Techniques for cloning the DNA encoding the protein of the present invention include, in addition to the methods described in the specification of the present application, PCR amplification using a synthetic DNA containing the nucleotide sequence of the present invention (e.g., SEQ ID NO: 2) as a primer, and selection of the DNA inserted into a suitable vector by hybridization with a labeled DNA fragment encoding a partial or full coding region of the protein of the present invention or a labeled synthetic DNA. Another technique involves direct amplification from total RNAs or mRNA fractions prepared from cells or tissues, using the reverse transcriptase polymerase chain reaction (RT-PCR method). As a DNA inserted into a suitable vector, for example, a commercially available library (e.g., from CLONTECH and STRATAGENE) can be used. Techniques for hybridization are normally used in the art, and can be easily carried out in accordance with various laboratory manuals such as T. Maniatis et al., supra. Depending on the intended purpose, the cloned DNA encoding the protein of the present invention can be used as such or if desired after digestion with a restriction enzyme or addition of a linker. The DNA thus obtained may be a gene having a nucleotide sequence of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 or 90, or a gene of above items (3) to (5). The DNA sequence to be inserted into an expression vector in the above step (B) may be a full-length cDNA encoding the above full-length protein or a DNA fragment, or a DNA fragment constructed so that it expresses a part thereof.

Thus, the present invention also includes a recombinant vector which comprises the above DNA sequence. The expression vector for

the protein of the present invention can be produced, for example, by excising the desired DNA fragment from the DNA encoding the protein of the present invention, and ligating the DNA fragment downstream of a promoter in a suitable expression vector.

Expression vectors for use in the present invention may be any vectors derived from prokaryotes (e.g., *E. coli*), yeast, fungi, insect viruses and vertebrate viruses so long as such vectors are replicable. However, the vectors which are compatible with microorganisms or cells used as hosts should be selected. Suitable combinations of host cell - expression vector systems are selected depending on the desired expression product.

When bacteria are used as hosts, plasmid vectors compatible with these bacteria are generally used as replicable expression vectors for recombinant DNA molecules.

For example, the plasmids pBR322 and pBR327 can be used to transform *E. coli*. Plasmid vectors normally contain an origin of replication, a promoter, and a marker gene conferring upon a recombinant DNA a phenotype useful for selecting the cells transformed with the recombinant DNA. Example of such promoters include a β -lactamase, lactose promoter and tryptophan promoter. Examples of such marker genes include an ampicillin resistance gene, and a tetracycline resistance gene. Examples of suitable expression vectors include the plasmids pUC18 and pUC19 in addition to pBR322, pBR327.

In order to express the DNA of the present invention in yeast, for example, YEp24 can be used as a replicable vector. The plasmid YEp24 contains the URA3 gene, which can be employed as a marker gene. Examples of promoters in expression vectors for yeast cells include promoters derived from genes for 3-phosphoglycerate kinase,

glyceraldehyde-3-phosphate dehydrogenase and alcohol dehydrogenase.

Examples of promoters and terminators for use in expression vectors to express the DNA of the present invention in fungal cells include promoters and terminators derived from genes for phosphoglycerate kinase (PGK), glyceraldehyde-3-phosphate dehydrogenase (GAPD) and actin. Examples of suitable expression vectors include the plasmids pPGACY2 and pBSFAHY83.

Examples of promoters for use in expression vectors to express the DNA of the present invention in insect cells include a polyhedrin promoter and P10 promoter.

Recombinant vectors used to express the DNA of the present invention in animal cells normally contain functional sequences to regulate genes, such as an origin of replication, a promoter to be placed upstream of the DNA of the present invention, a ribosome-binding site, a polyadenylation site and a transcription termination sequence. Such functional sequences which can be used to express the DNA of the present invention in eukaryotic cells can be obtained from viruses and viral substances. Examples of such functional sequences include an SR α promoter, SV40 promoter, LTR promoter, CMV (cytomegalovirus) promoter and HSV-TK promoter. Among them, a CMV promoter and SR α promoter can be preferably used. As promoters to be placed inherently upstream of the gene encoding the protein of the present invention, any promoters can be used so long as they are suitable for use in the above host-vector systems. Examples of origins of replication include foreign origins of replication, for example, those derived from viruses such as adenovirus, polyoma virus and SV40 virus. When vectors capable of integration into host chromosomes are used as expression vectors,

origins of replication of the host chromosomes may be employed. Examples of suitable expression vectors include the plasmids pSV-dhfr (ATCC 37146), pBPV-1(9-1) (ATCC 37111), pcDNA3.1 (INVITROGEN) and pME18S-FL3.

The present invention also includes a transformant which comprises the above recombinant vector.

Microorganisms or cells transformed with the replicable recombinant vector of the present invention can be selected from remaining untransformed parent cells based on at least one phenotype conferred by the recombinant vector. Phenotypes can be conferred by inserting at least one marker gene into the recombinant vector. Marker genes naturally contained in replicable vectors can be employed. Examples of marker genes include drug resistance genes such as neomycin resistance genes, and genes encoding dihydrofolate reductase.

As hosts for use in the above step (C), any of prokaryotes (e.g., *E. coli*), microorganisms (e.g., yeast and fungi) as well as insect and animal cells can be used so long as such hosts are compatible with the expression vectors used. Examples of such microorganisms include *Escherichia coli* strains such as *E. coli* K12 strain 294 (ATCC 31446), *E. coli* X1776 (ATCC 31537), *E. coli* C600, *E. coli* JM109 and *E. coli* B strain; bacterial strains belonging to the genus *Bacillus* such as *Bacillus subtilis*; intestinal bacteria other than *E. coli*, such as *Salmonella typhimurium* or *Serratia marcescens*; and various strains belonging to the genus *Pseudomonas*. Examples of such yeast include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Pichia pastoris*. Examples of such fungi include *Aspergillus nidulans*, and *Acremonium chrysogenum* (ATCC 11550).

As insect cells, for example, *Spodoptera frugiperda* (Sf cells), High Five™ cells derived from eggs of *Trichoplusia ni*, etc., can be used when the virus is AcNPV. Examples of such animal cells include HEK 293 cells, COS-1 cells, COS-7 cells, Hela cells, and Chinese hamster ovary (CHO) cells. Among them, CHO cells, HEK 293 cells, etc., are preferred.

When cells are used as hosts, combinations of expression vectors and host cells to be used vary with experimental objects. According to such combinations, two types of expression (i.e. transient expression and constitutive expression) can be included.

"Transformation" of microorganisms and cells in the above step (C) refers to introducing DNA into microorganisms or cells by forcible methods or phagocytosis of cells and then transiently or constitutively expressing the trait of the DNA in a plasmid or an intra-chromosome integrated form. Those skilled in the art can carry out transformation by known methods [see e.g., "Idenshi Kougaku Handbook (Genetic Engineering Handbook)", an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.]. For example, in the case of animal cells, DNA can be introduced into cells by known methods such as DEAE-dextran method, calcium-phosphate-mediated transfection, electroporation, lipofection, etc. For stable expression of the protein of the present invention using animal cells, there is a method in which selection can be carried out by clonal selection of the animal cells containing the chromosomes into which the introduced expression vectors have been integrated. For example, transformants can be selected using the above selectable marker as an indication of successful transformation. In addition, the animal cells thus obtained using the selectable marker can be subjected to repeated clonal selection

to obtain stable animal cell strains highly capable of expressing the protein of the present invention. When a dihydrofolate reductase (DHFR) gene is used as a selectable marker, one can culture animal cells while gradually increasing the concentration of methotrexate (MTX) and select the resistant strains, thereby amplifying the DNA encoding the protein of the present invention together with the DHFR gene to obtain animal cell strains having higher levels of expression.

The above transformant can be cultured under conditions which permit the expression of the DNA encoding the protein of the present invention to produce and accumulate the protein of the present invention. In this manner, the protein of the present invention or a salt thereof can be produced. Thus, the present invention also includes a process for producing a protein, which comprises expressing the protein encoded by the above DNA from the above transformant.

The above transformant can be cultured by methods known to those skilled in the art (see e.g., "Bio Manual Series 4", YODOSHA CO., LTD.). For example, animal cells can be cultured by various known animal cell culture methods including attachment culture such as Petri dish culture, multitray type culture and module culture, attachment culture in which cells are attached to cell culture carriers (microcarriers), suspension culture in which productive cells themselves are suspended. Examples of media for use in the culture include media commonly used for animal cell culture, such as D-MEM and RPMI 1640.

In order to separate and purify the protein of the present invention from the above culture, suitable combinations of per se known separation and purification methods can be used. Examples

such methods include methods based on solubility, such as salting-out and solvent precipitation; methods based on the difference in charges, such as ion-exchange chromatography; methods mainly based on the difference in molecular weights, such as dialysis, ultrafiltration, gel filtration and SDS-polyacrylamide gel electrophoresis; methods based on specific affinity, such as affinity chromatography; methods based on the difference in hydrophobicity, such as reverse phase high performance liquid chromatography; and methods based on the difference in isoelectric points, such as isoelectric focusing.

The protein of the present invention can also be produced as a fusion protein with another protein. These fusion proteins are also included within the present invention. For the expression of such fusion proteins, any vectors can be used so long as the DNA encoding the protein can be inserted into the vectors and the vectors can express the fusion protein. Examples of fusion protein include glutathione S-transferase (GST) and a hexa-histidine sequence (6 x His). The fusion protein of the protein of the present invention with another protein can be advantageously purified by affinity chromatography using a substance with an affinity for the fusion partner protein. For example, fusion proteins with GST can be purified by affinity chromatography using glutathione as a ligand.

The present invention also includes an inhibitory protein, i.e., a protein capable of inhibiting the activity of the protein of above item (6). Examples of such inhibitory proteins include antibodies, or proteins which bind to active sites of proteins, thereby inhibiting the expression of their activity.

The present invention also relates to an antibody directed against the protein of the present invention or a fragment thereof,

and to its production. The antibody is not specifically limited so long as it can recognize the protein of the present invention. Examples of such antibodies include polyclonal antibodies, monoclonal antibodies and their fragments, single chain antibodies and humanized antibodies. Antibody fragments can be produced by known techniques. Examples of such antibody fragments include, but not limited to, $F(ab')_2$ fragments, Fab' fragments, Fab fragments and Fv fragments. The antibody directed against the protein of the present invention can be produced using the protein of the present invention or a peptide thereof as an immunogen according to per se known process for producing antibodies or antisera. Such methods are described, for example, in "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", the third edition, an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.

In the case of polyclonal antibodies, for example, the protein of the present invention or a peptide thereof can be injected to animals such as rabbits to produce antibodies directed against the protein or peptide, and then their blood can be collected. The polyclonal antibodies can be purified from the blood, for example, by ammonium sulfate precipitation or ion-exchange chromatography, or by using the affinity column on which the protein has been immobilized.

In the case of monoclonal antibodies, for example, animals such as mice are immunized with the protein of the present invention, their spleen is removed and homogenized to obtain spleen cells, which are then fused with mouse myeloma cells by using a reagent such as polyethylene glycol. From the resulting hybrid cells (i.e. hybridoma cells), the clone producing the antibody directed against the protein of the present invention can be selected. Then, the

resulting clonal hybridoma cells can be implanted intraperitoneally into mice, the ascitic fluid recovered from the mice. The resulting monoclonal antibody can be purified, for example, by ammonium sulfate precipitation or ion-exchange chromatography, or by using the affinity column on which the protein has been immobilized.

When the resulting antibody is used to administer it to humans, it is preferably used as a humanized antibody or human antibody in order to reduce its immunogenicity. The humanized antibody can be produced using transgenic mice or other mammals. For a general review of humanized antibodies, see, for example, Morrison, S.L. et al., Proc. Natl. Acad. Sci. USA, 81:6851-6855 (1984); Jones, P.T. et al., Nature 321:522-525 (1986); Hiroshi Noguchi, Igaku no Ayumi (J. Clin. Exp. Med.) 167:457-462 (1993); Takashi Matsumoto, Kagaku to Seibutsu (Chemistry and Biology) 36:448-456 (1998). Humanized chimeric antibodies can be produced by linking a V region of a mouse antibody to a C region of a human antibody. Humanized antibodies can be produced by substituting a sequence derived from a human antibody for a region other than a complementarity-determining region from a mouse monoclonal antibody. In addition, human antibodies can be directly produced in the same manner as the production of conventional monoclonal antibodies by immunizing the mice whose immune systems have been replaced with human immune systems. These antibodies can be used to purify the protein of the present invention from a cell extract or transformed cells producing the protein of the present invention. These proteins can also be used to construct ELISA, RIA (radioimmunoassay) and western blotting systems. These assay systems can be used for diagnostic purposes. These antibodies can also be used as reagents for studying functions of the protein of the present invention.

The antibodies of the present invention can be purified and then administered to patients. Thus in another aspect, the present invention is a pharmaceutical composition which comprises the above antibody as an active ingredient, and therapy using the antibody of the present invention. In such pharmaceutical compositions, the active ingredient may be combined with other therapeutically active ingredients or inactive ingredients (e.g., conventional pharmaceutically acceptable carriers or diluents such as immunogenic adjuvants) and physiologically non-toxic stabilizers and excipients. The resulting combinations can be sterilized by filtration, and formulated into vials after lyophilization or into various dosage forms in stabilized and preservable aqueous preparations. These antibodies exhibit therapeutic activity by inhibiting the NF- κ B activation mediated by the protein of the present invention.

The DNA of the present invention can also be used to isolate, identify and clone other proteins involved in intracellular signal transduction processes. For example, the DNA sequence encoding the protein of the present invention can be used as a "bait" in yeast two-hybrid systems (see e.g., Nature 340:245-246 (1989)) to isolate and clone the sequence encoding a protein ("prey") which can associate with the protein of the present invention. In a similar manner, it can be determined whether the protein of the present invention can associate with other cellular proteins (e.g., NIK and TRAF2). In another method, proteins which can associate with the protein of the present invention can be isolated from cell extracts by immunoprecipitation [see e.g., "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.] using antibodies

directed against the protein of the present invention. In still another method, the protein of the present invention can be expressed as a fusion protein with another protein as described above, and immunoprecipitated with an antibody directed against the fusion protein in order to isolate a protein which can associate with the protein of the present invention.

The DNA of the present invention can be used to detect abnormality in the DNA or mRNA encoding the protein of the present invention or a peptide fragment thereof. Thus, for example, the DNA of the present invention is useful for gene diagnosis regarding damage, mutations, and reduced, increased or over-expression of the DNA or mRNA. That is, the present invention includes a method for diagnosing a disease associated with the expression or activity of NF- κ B in a subject, which comprises the steps of:

- (a) determining the presence or absence of a mutation in the NF- κ B coding nucleotide sequence in the genome of the subject; and/or
- (b) measuring the amount of the expression of the protein according to above item (6) in the subject, and comparing it with a normal value.

When the NF- κ B coding nucleotide sequence contains a mutation according to the above step (a), the mutation may cause disease associated with the expression or activity of NF- κ B. When the amount of the expression of the protein of above item (6) is different from the normal value according to the above step (b), the abnormal expression of the NF- κ B activating new protein of the present invention may be responsible for diseases associated with the expression or activity of NF- κ B. Determination of the presence or absence of a mutation in the NF- κ B coding nucleotide sequence in the above step (a) may involve RT-PCR using a part of the

nucleotide sequence of the NF- κ B gene as a primer, followed by conventional DNA sequencing to detect the presence or absence of the mutation. PCR-SSCP [Genomics 5:874-879 (1989); "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.] can also be used to determine the presence or absence of the mutation. Measurement of the amount of the expression of the protein in the above step (b) may involve, for example, using the antibody of above item (15). Sequencing the nucleotide sequence may involve, for example, RT-PCR using a part of the nucleotide sequence of the gene of above items (3) to (5) as a primer, followed by conventional DNA sequencing to detect the presence or absence of the mutation. PCR-SSCP [Genomics 5:874-879 (1989); "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.] can also be used to determine the presence or absence of the mutation.

The present invention also relates to a method for screening for a compound which inhibits or activates the activation of the protein NF- κ B of the present invention. Such method comprises the steps of (a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal; (b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell; (c) contacting the transformed cell with one or more candidate compounds; and (d) measuring the detectable signal. Examples of components capable of providing a detectable signal include reporter genes. Reporter genes are used instead of directly detecting the activation of transcription factors of interest. The transcriptional activity of a promoter of a gene is analyzed by

linking the promoter to a reporter gene and measuring the activity of the product of the reporter gene ("Bio Manual Series 4" (1994), YODOSHA CO., LTD.). Any peptide or protein can be used so long as those skilled in the art can measure the activity or amount of the expression product (including the amount of the produced mRNA) of the reporter genes. For example, enzymatic activity of chloramphenicol acetyltransferase, β -galactosidase, luciferase, etc., can be measured. Any reporter plasmids can be used to evaluate NF- κ B activation so long as the reporter plasmids have an NF- κ B recognition sequence inserted upstream of the reporter gene. For example, pNF- κ B-Luc (STRATAGEGE) can be used. Other examples include NF- κ B dependent reporter plasmids described in Tanaka S. et al., J. Vet. Med. Sci. Vol.59 (7); Rothe M. et al., Science Vol.269, p.1424-1427 (1995). Any host cells can be used so long as NF- κ B activation can be detected in the host cells. Preferred host cells are mammal cells such as 293-EBNA cells. Transformation and culture of the cells can be carried out as described above. In a specific embodiment, the method for screening a compound which inhibits or activates NF- κ B activation comprises culturing the transformed cell for a certain period of time, adding a certain amount of a test compound, measuring the reporter activity expressed by the cell after a certain period of time, and comparing the activity with that of a cell to which the test compound has not been added. The reporter activity can be measured by methods known in the art (see e.g., "Bio Manual Series 4" (1994), YODOSHA CO., LTD.). Examples of test compounds include, but not limited to, low molecular weight compounds and peptides. Test compounds may be artificially synthesized compounds or naturally occurring compounds. Test compounds may be a single compound or mixtures. Examples of such

detectable signals which may be measured include the amount of mRNA or proteins for genes whose expression is known to be induced by NF- κ B activation (e.g., genes for IL-1 and TNF- α , in addition to the above reporter genes).

Thus, compounds which inhibit or activate NF- κ B activation may be screened by the steps of:

- (a) transforming a host cell with a gene encoding a protein capable of activating NF- κ B;
- (b) culturing the transformed cell under conditions which permit the expression of the gene in the host cell;
- (c) contacting the transformed cell with one or more candidate compounds; and
- (d) measuring the amount of NF- κ B or mRNA of a protein whose expression is induced by NF- κ B, or the amount of such protein.

The amount of mRNA can be measured, for example, by northern hybridization, RT-PCR, etc. The amount of proteins can be measured, for example, by using antibodies. The antibodies may be produced by known methods. Commercially available antibodies (from, e.g., Wako Pure Chemical Industries, Ltd.) can also be used.

The present invention also includes a compound obtainable by the above screening method. However, the screening method of the present invention is not limited to the above method. The present invention also includes a process for producing the above compound by the method of above item (12).

The present invention also includes a kit for screening a compound which inhibits or activates NF- κ B activation. The kit comprises (a) a host cell transformed with a gene encoding a protein capable of activating NF- κ B, and a component capable of providing a detectable signal; and (b) a medium which permits the expression

of the gene in the host cell. Thus , the present invention provides the kit which contains reagents necessary for such screening.

Because NF- κ B is involved in a wide variety of pathological conditions such as inflammation, autoimmune diseases, cancers and viral infections, it is an attractive target for drug design and therapeutic intervention. Many experiments show that the inhibition of NF- κ B activity may have significant physiological effects [see e.g., Ann. Rheum. Dis. 57:738-741 (1998); American Journal of Pathology 152:793-803 (1998); ARTHRITIS & RHEUMATISM 40:226-236 (1997); Am. J. Respir. Crit. Care Med. 158:1585-1592 (1998); J. Exp. Med. 188:1739-1750 (1998); Gut 42:477-484 (1998); The Journal of Immunology 161:4572-4582 (1998); Nature Medicine 3:894-899 (1997)]. The finding of the new protein described herein capable of activating NF- κ B has provided a new method for inhibiting an abnormal NF- κ B function. Thus, the present invention also relates to use of a compound which inhibits the function of the protein capable of activating NF- κ B described above, for inhibiting NF- κ B activation. The compound obtained by the above screening method, which inhibits NF- κ B activation, is useful as a medicament to treat or prevent diseases characterized by undesirable activation of NF- κ B, such as inflammation, autoimmune diseases, infectious diseases (e.g., HIV infection) and cancers. Recently, it has also become apparent that NF- κ B activation inhibits apoptosis of cells. The compound obtained by the above screening method, which inhibits NF- κ B activation, may be capable of stimulating apoptosis. Diseases which may be treated by the induction of apoptosis include tumors.

On the other hand, diseases which may be treated by the inhibition of apoptosis include GVHD, skin diseases such as toxic

epidermal necrolysis (TEN), proliferative nephritides (e.g., IgA nephritis, purpuric nephritis and lupus nephritis) and fulminant hepatitis. Thus, the compound obtained by the above screening method, which activates NF- κ B activation, is useful as a medicament to treat or prevent these diseases.

In addition, the gene encoding the protein of the present invention is useful for gene therapy to treat various diseases such as cancers, autoimmune diseases, allergy diseases and inflammatory response. "Gene therapy" refers to administering into the human body a gene or a cell into which a gene has been introduced. The protein of the present invention and the DNA encoding the protein can also be used for diagnostic purposes.

The compound obtained by the screening method of the present invention or a salt thereof can be formulated into the above pharmaceutical compositions (e.g., tablets, capsules, elixirs, microcapsules, sterile solutions and suspensions) according to conventional procedures. The formulations thus obtained are safe and low toxic, and can be administered, for example, to humans and mammals (e.g., rats, rabbits, sheep, pigs, cattle, cats, dogs and monkeys). Administration to patients can be carried out by methods known in the art, such as intra-arterial injection, intravenous injection and subcutaneous injection. The dosage may vary with the weight and age of the patient as well as a mode of administration, but those skilled in the art can appropriately select suitable dosages. When the compound can be encoded by DNA, the DNA can be inserted into a vector for gene therapy, and gene therapy can be carried out. The dosage and mode of administration may vary with the weight, age and symptoms of the patient, but those skilled in the art can appropriately select them. Thus, the present invention

also relates to a pharmaceutical composition which comprises the above compound as an active ingredient.

In addition, the above compound is useful as a medicament to treat or prevent diseases characterized by undesirable activation of NF- κ B, such as inflammation, autoimmune diseases, viral diseases, infectious diseases and cancers. Thus, the present invention also relates to a pharmaceutical composition for inflammation, autoimmune diseases, viral diseases, cancers, etc., which comprises the above compound. Specifically, the pharmaceutical composition is useful as a therapeutic and prophylactic drug against, for example, rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, diabetes, sepsis, asthma, allergic rhinitis, ischemic heart diseases, inflammatory intestinal diseases, subarachnoid hemorrhage, viral hepatitis and AIDS.

The present invention also relates to the use of a compound of above item (13) for manufacturing a medicament against inflammation, autoimmune diseases, viral diseases, cancers, etc. The present invention also includes an antisense oligonucleotide against a gene of any one of above items (3) to (5). An antisense oligonucleotide refers to an oligonucleotide complementary to the target gene sequence. The antisense oligonucleotide can inhibit the expression of the target gene by inhibiting RNA functions such as translation to proteins, transport to the cytoplasm and other activity necessary for overall biological functions. In this case, the antisense oligonucleotide may be RNA or DNA. The DNA sequence of the present invention can be used to produce an antisense oligonucleotide capable of hybridizing with the mRNA transcribed from the gene encoding the protein of the present invention. It is known that an antisense oligonucleotide generally has an

inhibitory effect on the expression of the corresponding gene (see e.g., Saibou Kougaku Vol.13, No.4 (1994)). The oligonucleotide containing an antisense coding sequence against a gene encoding the protein of the present invention can be introduced into a cell by standard methods. The oligonucleotide effectively blocks the translation of mRNA of the gene encoding the protein of the present invention, thereby blocking its expression and inhibiting undesirable activity.

The oligonucleotide of the present invention may be a naturally occurring oligonucleotide or its modified form [see e.g., Murakami & Makino, Saibou Kougaku Vol.13, No.4, p.259-266 (1994); Akira Murakami, Tanpakushitsu Kakusan Kouso (PROTEIN, NUCLEIC ACID AND ENZYME) Vol.40, No.10, p.1364-1370 (1995)]. Thus, the oligonucleotide may have modified sugar moieties or inter-sugar moieties. Examples of such modified forms include phosphothioates and other sulfur-containing species used in the art. According to several preferred embodiments of the present invention, at least one phosphodiester bond in the oligonucleotide is substituted with the structure which can enhance the ability of the composition to permeate cellular regions where RNA with the activity to be regulated is located.

Such substitution preferably involves a phosphorothioate bond, a phosphoramidate bond, methylphosphonate bond, or a short-chain alkyl or cycloalkyl structure. The oligonucleotide may also contain at least some modified base forms. Thus, it may contain purine and pyrimidine derivatives other than naturally occurring purine and pyrimidine. Similarly, the furanosyl moieties of the nucleotide subunits can be modified so long as the essential purpose of the present invention is attained. Examples of such

modifications include 2'-O-alkyl and 2'-halogen substituted nucleotides. Examples of modifications in sugar moieties at their 2-position include OH, SH, SCH₃, OCH₃, OCN or O(CH₂)_nCH₃, wherein n is 1 to about 10, and other substituents having similar properties. All the analogues are included in the scope of the present invention so long as they can hybridize with the mRNA of the gene of the present invention to inhibit functions of the mRNA.

The oligonucleotide of the present invention contains about 3 to about 50 nucleotides, preferably about 8 to about 25 nucleotides, more preferably about 12 to about 20 nucleotides. The oligonucleotide of the present invention can be produced by the well-known solid phase synthesis technique. Devices for such synthesis are commercially available from some manufactures including Applied Biosystems. Other oligonucleotides such as phosphothioates can also be produced by methods known in the art.

The oligonucleotide of the present invention is designed to hybridize with the mRNA transcribed from the gene of the present invention. Those skilled in the art can easily design an antisense oligonucleotides based on a given gene sequence.

Another method includes the triple helix technique. This technique involves forming a triple helix on the targeted intra-nuclear DNA sequence, thereby regulating its gene expression, mainly at the transcription stage. The oligonucleotide is designed mainly in the gene region involved in the transcription and inhibits the transcription and the production of the protein of the present invention. Such RNA, DNA and oligonucleotide can be produced using known synthesizers.

The oligonucleotide may be introduced into the cells containing the target nucleic acid sequence by any of DNA

transfection methods such as calcium phosphate method, electroporation, lipofection, microinjection, or gene transfer methods including the use of gene transfer vectors such as viruses. An antisense oligonucleotide expression vector can be prepared using a suitable retrovirus vector, then the expression vector can be introduced into the cells containing the target nucleic acid sequence by contacting the vector with the cells in vivo or ex vivo.

The DNA of the present invention can be used in the antisense RNA/DNA technique or the triple helix technique to inhibit NF- κ B activation mediated by the protein of the present invention.

The antisense oligonucleotide against the gene encoding the protein of the present invention is useful as a medicament to treat or prevent diseases characterized by undesirable activation of NF- κ B, such as inflammation, autoimmune diseases, infectious diseases (e.g., HIV infection) and cancers. Thus, the present invention also includes a pharmaceutical composition which comprises the above antisense oligonucleotide as an active ingredient. The antisense oligonucleotide can also be used to detect such diseases using northern hybridization or PCR.

The present invention also includes a ribozyme which inhibits NF- κ B activation. A ribozyme is an RNA capable of recognizing a nucleotide sequence of a nucleic acid and cleaving the nucleic acid (see e.g., Hiroshi Yanagawa, "Jikken Igaku (Experimental Medicine) Bioscience 12: New Age of RNA). The ribozyme can be produced so that it cleaves the selected target RNA (e.g., mRNA encoding the protein of the present invention). Based on the nucleotide sequence of the DNA encoding the protein of the present invention, the ribozyme specifically cleaving the mRNA of the protein of the present invention can be designed. Such ribozyme has a complementary

sequence to the mRNA for the protein of the present invention, complementarily associates with the mRNA and then cleaves the mRNA, which results in reduction or entire loss of the expression of the protein of the present invention. The level of the reduction of the expression is dependent on the level of the ribozyme expression in the target cells.

There are two types of ribozyme commonly used: a hammerhead ribozyme and a hairpin ribozyme. In particular, hammerhead ribozymes have been well studied regarding their primary and secondary structure necessary for their cleavage activity, and those skilled in the art can easily design the ribozymes nucleotided solely on the nucleotide sequence information for the DNA encoding the protein of the present invention [see e.g., Iida et al., Saibou Kougaku Vol.16, No.3, p.438-445 (1997); Ohkawa & Taira, Jikken Igaku (Experimental Medicine) Vol.12, No.12, p.83-88 (1994)]. It is known that the hammerhead ribozymes have a structure consisting of two recognition sites (recognition site I and recognition site II forming a chain complementary to target RNA) and an active site, and cleave the target RNA at the 3'end of its sequence NUX (wherein N is A or G or C or U, and X is A or C or U) after the formation of a complementary pair with the target RNA in the recognition sites. In particular, the sequence GUC (or GUA) has been found to have the highest activity [see e.g., Koizumi, M. et al., Nucl. Acids Res. 17:7059-7071 (1989); Iida et al., Saibou Kougaku Vol.16, No.3, p.438-445 (1997); Ohkawa & Taira, Jikken Igaku (Experimental Medicine) Vol.12, No.12, p.83-88 (1994); Kawasaki & Taira, Jikken Igaku (Experimental Medicine) Vol.18, No.3, p.381-386 (2000)].

Therefore the sequence GTC (or GTA) is searched out, and a ribozyme is designed to form several to ten-odd complementary base

pairs around that sequence. The suitability of the designed ribozyme can be evaluated by checking whether the prepared ribozyme can cleave the target mRNA in vitro. The ribozyme can be prepared by methods known in the art to synthesize RNA molecules.

Alternatively, the sequence of the ribozyme can be synthesized on a DNA synthesizer and inserted into various vectors containing a suitable RNA polymerase promoter (e.g., T7 or SP6) to enzymatically synthesize an RNA molecule in vitro. Such ribozymes can be introduced into cells by gene transfer methods such as microinjection. Another method involves inserting a ribozyme DNA into a suitable expression vector and introducing the vector into cell strains, cells or tissues. Suitable vectors can be used to introduce the ribozyme into a selected cell. Examples of vectors commonly used for such purpose include plasmid vectors and animal virus vectors (e.g., retrovirus, adenovirus, herpes or vaccinia virus vectors). Such ribozymes are capable of inhibiting the NF- κ B activation mediated by the protein of the present invention.

The present invention also relates to a process for obtaining a new gene having a function, which comprises using the oligo-capping method to construct a full-length cDNA library, and using a reporter gene indicative of the presence of a protein having the function.

Methods using a cDNA library containing a lot of non-full-length cDNAs are inefficient in obtaining many genes (cDNAs) having functions. Therefore libraries with a high ratio of the number of the full-length cDNA clones to the total number of the clones are necessary. "Full-length cDNA" refers to a complete DNA copy of mRNA from a gene. The cDNA libraries produced using the oligo-capping method contain full-length cDNA clones in a ratio of 50 to 80%, namely, a 5 to 10-fold increase in full-length cDNA clones

compared to the cDNA libraries produced by prior art methods (Sumio Sugano, the monthly magazine BIO INDUSTRY Vol.16, No.11, p.19-26). Full-length cDNA clones are essential for protein expression in functional analyses of genes, and full-length cDNA clones themselves are very important materials for activity measurement. Thus, cloning of full-length cDNA is necessary for functional analyses of genes. Sequencing of the cDNA not only provides important information for establishing the primary sequence of the protein encoded by the cDNA, but also reveals the entire exon sequence. Thus, the full-length cDNA provides valuable information for identifying a gene, such as information for determining the primary sequence of a protein, exon-intron structure, the transcription initiation site of mRNA, the location of a promoter, etc.

The construction of full-length cDNA libraries by the oligo-capping method can be carried out, for example, according to the method described in "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", the third edition (1999), an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD. The reporter gene indicative of the presence of a protein having a function contains one or more suitable expression regulation sequence portion to which a protein factor such as a transcriptional factor can bind, and a structural gene portion which allows the measurement of the activation of the proteins factor. The structural gene portion may encode any peptide or protein so long as those skilled in the art can measure the activity or amount of its expression product (including the amount of the mRNA produced). For example, chloramphenicol acetyltransferase, β -galactosidase, luciferase, etc., can be used and their enzymatic activity measured.

The oligo-capping method used herein involves substituting a cap structure with a synthetic oligo sequence by using BAP, TAP and an RNA ligase, as described in Suzuki & Sugano, "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", the third edition (1999), an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.

The process of the present invention uses an in vitro system or a cell-based system, preferably a cell-based system. Examples of such cells include cells of prokaryotes such as E. coli, microorganisms such as yeast and fungi, as well as insects and animals. Preferred examples include animal cells, in particular, 293-EBNA cells and NIH3T3 cells.

Examples of reporter genes indicative of the presence of a protein having a function include reporter genes containing a CREB (cAMP responsive element binding protein) binding sequence or AP-1 (activator protein-1) binding sequence at the expression regulation sequence region of the reporter genes, in addition to the NF- κ B reporter genes described herein. For example, if a gene capable of activating CREB is to be obtained, a CREB-dependent reporter plasmid and a full-length cDNA clone produced by the oligo-capping method can be cotransfected into cells, and a plasmid having increased reporter activity can be selected from the cells to attain the purpose. If a gene capable of inhibiting CREB is to be obtained, a CREB-dependent reporter plasmid and a full-length cDNA clone produced by the oligo-capping method can be cotransfected into cells, and a plasmid having decreased reporter activity can be selected from the cells to attain the purpose. These procedures may be carried out in the presence of a certain stimulus to the cells. The cDNA to be transfected into the cells may be a single clone or

multiple clones which may be transfected simultaneously. One embodiment of the process of the present invention is detailed in Examples herein. Alternatively, a screening system for obtaining a gene capable of inhibiting NF- κ B activation can also be constructed by cotransfecting a full-length cDNA and a reporter gene into cells and selecting a clone having subnormally increased reporter activity.

However, the process of the present invention is not limited to these embodiments.

Because the cDNA of the present invention is full-length, its 5' end sequence is the transcription initiation site of the corresponding mRNA. Therefore the cDNA sequence can be used to identify the promoter region of the gene by comparing the cDNA with the genomic nucleotide sequence. Genomic nucleotide sequences are available from various databases when the sequences have been deposited in the databases. Alternatively, the cDNA can also be used to clone the desired sequence from a genomic library, for example, by hybridization, and determine its nucleotide sequence. Thus, by comparing the nucleotide sequence of the cDNA of the present invention with a genomic sequence, the promoter region of the gene located upstream the cDNA can be identified. In addition, the promoter fragment thus identified can be used to construct a reporter plasmid for evaluating the expression of the gene. In general, the DNA fragment spanning 2kb (preferably 1kb) upstream from the transcription initiation site can be inserted upstream of the reporter gene to produce the reporter plasmid. The reporter plasmid can be used to screen for a compound which enhances or reduces the expression of the gene. For example, such screening can be carried out by transforming a suitable cell with the reporter plasmid,

culturing the transformed cell for a certain period of time, adding a certain amount of a test compound, measuring the reporter activity expressed by the cell after a certain period of time, and comparing the activity with that of a cell to which the test compound has not been added. These methods are also included in the scope of the present invention.

The present invention also relates to a computer-readable medium on which a sequence data set has been stored, said sequence data set comprising a nucleotide sequence expressed by SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90.

In another aspect, the present invention relates to a method for calculating a homology, which comprises comparing data on the above medium with data of other nucleotide sequences. Thus, the gene and amino acid sequence of the present invention provide valuable information for determining their secondary and tertiary structure, e.g., information for identifying other sequence having a similar function and high homology. These sequences are stored on the computer-readable medium, then a database is searched using data stored in a known macromolecule structure program and a known search tool such as GCG. In this manner, a sequence in a database having a certain homology can be easily found.

The computer-readable medium may be any composition of materials used to store information or data. Examples of such media include commercially available floppy disks, tapes, chips, hard drives, compact disks and video disks. The data on the medium allows a method for calculating a homology by comparing the data with other nucleotide sequence data. This method comprises the steps of

providing a first polynucleotide sequence containing the polynucleotide sequence of the present invention for the computer-readable medium, and then comparing the first polynucleotide sequence with at least one-second polynucleotide or polypeptide sequence to identify the homology.

Examples:

The following examples further illustrate, but do not limit the present invention.

Example 1: Construction of a full-length cDNA library using the oligo-capping method

(1) Preparation of RNA from human lung fibroblasts (Cryo NHLF)

Human lung fibroblasts (Cryo NHLF: purchased from Sanko Junyaku Co., Ltd.) were cultured according to the attached protocol. After repeating subculturing the cells to obtain fifty 10cm dishes containing the resulting culture, the cells were recovered with a cell scraper. Then, total RNA was obtained from the recovered cells by using the RNA extraction reagent ISOGEN (purchased from NIPPON GENE) according to the manufacture's protocol. Then, poly A⁺ RNA was obtained from the total RNA by using an oligo-dT cellulose column according to Maniatis et al., supra.

(2) Construction of a full-length cDNA library by the oligo-capping method

A full-length cDNA library was constructed from the above poly A⁺ RNA by the oligo-capping method according to the method of Sugano S. et al. [e.g., Maruyama, K. & Sugano, S., Gene, 138:171-174 (1994); Suzuki, Y. et al., Gene, 200:149-156 (1997); Suzuki, Y. & Sugano, S. "Shin Idenshi Kougaku Handbook (New Genetic Engineering Handbook)", the third edition (1999), an extra issue of "Jikken Igaku (Experimental Medicine)", YODOSHA CO., LTD.].

(3) Preparation of plasmid DNA

The full-length cDNA library constructed as above was transfected into E. coli strain TOP 10 by electroporation, then spread on LB agar medium, and incubated overnight at 37°C. Then, using QIAwell 96 Ultra Plasmid Kit (QIAGEN) according to the attached protocol, the plasmids were recovered from the colonies grown on ampicillin-containing LB agar medium.

Example 2: Cloning of DNA capable of activating NF- κ B

(1) Screening of the cDNA encoding the protein capable of activating NF- κ B

293-EBNA cells (purchased from Invitrogen) were grown to 1×10^4 cells/well for 24 hours at 37°C (in the presence of 5% CO₂) using 5% FBS containing DMEM medium. Then, 50ng of pNF κ B-Luc (purchased from STRATAGENE) and 2 μ l of the full-length cDNA prepared in above Example 1.(3) were cotransfected into the cells in a well using FuGENE 6 (purchased from Roche) according to the attached protocol. After 24 hours of culture at 37°C, the reporter activity of NF- κ B was measured using PIKKA GENE LT2.0 (TOYO INK). The luciferase activity was measured using Wallac ARVO™ST 1420 MULTILABEL COUNTER (Perkin Elmer).

(2) DNA sequencing

The above screening was carried out for 115,000 clones, and plasmids showing a 10-fold or more increase in luciferase activity compared to that of the control experiment were selected. One pass sequencing was carried out from the 5' end of the cloned cDNA (sequencing primer: 5'-CTTCTGCTCTAAAAGCTGCG-3' (SEQ ID NO: 3)) and from the 3' end (sequencing primer: 5'-CGACCTGCAGCTCGAGCACA-3' (SEQ ID NO: 4)) so that as long sequence as possible is determined. The sequencing was carried out using the reagent Thermo Sequenase II

Dye Terminator Cycle Sequencing Kit (Amersham Pharmacia Biotech) or BigDye Terminator Cycle Sequencing FS Ready Reaction Kit (Applied Biosystems) and the device ABI PRISM 377 sequencer or ABI PRISM 3100 sequencer according to the attached instructions.

(3) Database analysis of the obtained clones

BLAST (Basic local alignment search tool) searching [S. F. Altschul et al., J. Mol. Biol., 215:403-410 (1990)] was carried out in GenBank for the obtained nucleotide sequence. The results showed that 75 clones represented 45 genes encoding new proteins capable of activating NF- κ B.

(4) Full-length sequencing

The full-length DNA sequences for the 45 new clones were determined (SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88 and 90). The amino acid sequences of the protein coding regions (open reading frames) were deduced (SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87 and 89).

Example 3: Screening compounds inhibiting NF- κ B activation

293-EBNA cells were seeded on 5% FBS containing DMEM medium in a 96-well cell culture plate to a final cell density of 1×10^4 cells/100 μ l/well, and cultured for 24 hours at 37°C in the presence of 5% CO₂. Then, 10ng of the plasmid containing the gene encoding NF- κ B activating protein of SEQ ID NO: 81 and 50ng of the reporter plasmid pNF κ B-Luc were cotransfected into the cells in a well using FuGENE 6. After 1 hour, the proteasome inhibitor MG132 (purchased from CALBIOCHEM) (Uehara T. et al., J. Biol. Chem. 274, p.15875-15882

(1999); Wang X. C. et al., Invest. Ophthalmol. Vis. Sci. 40, p.477-486) was added to the culture to a final concentration of 10 μ M. After 24 hours of culture at 37°C, the reporter activity was measured using PIKKA GENE LT2.0. The results showed that MG132 inhibited the expression of the reporter gene (Fig. 1).

EFFECTS OF THE INVENTION

As described above, the present invention provides industrially highly useful proteins capable of activating NF- κ B and genes encoding the proteins. The proteins of the present invention and the genes encoding the proteins allow not only screening for compounds useful for treating and preventing diseases associated with the excessive activation or inhibition of NF- κ B, but also production of diagnostics for such diseases. The genes of the present invention are also useful as a gene source used for gene therapy.

FREE TEXT OF SEQUENCE LISTING

SEQ ID NO: 91 and SEQ ID NO: 92 are primers.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph showing reporter activity inhibition by the proteasome inhibitor MG132 in Example 3, the axis of abscissa is M132 concentration and the transversal axis is relative luciferase activity.

NAME OF THE DOCUMENT: ABSTRACT

ABSTRACT

PROBLEMS TO BE SOLVED: Provision of proteins having NF- κ B activity, which are used for diagnosing, treating or preventing diseases associated with the excessive activation or inhibition of NF- κ B.

MEANS TO SOLVE THE PROBLEMS: Using the plasmid pNF κ B-Luc, the cDNA encoding a protein capable of activating NF- κ B has been cloned from the cDNA library constructed from human lung fibroblasts, and the DNA sequence and the deduced amino acid sequence determined. The protein, the DNA encoding the protein, a recombinant vector containing the DNA, and a transformant containing the recombinant vector are useful for screening a substance inhibiting or activating NF- κ B activation.

FIG.1

